

AD-A040 202

NAVAL POSTGRADUATE SCHOOL MONTEREY CALIF
A DIFFUSION APPROXIMATION ANALYSIS OF A GENERAL N-COMPARTMENT S--ETC(U)
APR 77 D P GAVER, J P LEHOCZKY
NPS55-77-16

F/G 12/1

NL

UNCLASSIFIED

| OF |

AD
A040 202



END

DATE
FILMED
6-77

AD A 040202

NPS55-77-16

NAVAL POSTGRADUATE SCHOOL

Monterey, California



AD No. —
DDC FILE COPY.

A DIFFUSION APPROXIMATION ANALYSIS
OF A GENERAL n -COMPARTMENT SYSTEM

by

Donald P. Gaver

and

John P. Lehoczký

April 1977

Approved for public release; distribution unlimited.

Research supported in part by Grant AFOSR 74-2642
from the Air Force Office of Scientific Research.

Research supported by NSF Grant AG467 at the Naval
Postgraduate School.

NAVAL POSTGRADUATE SCHOOL
Monterey, California

Rear Admiral Isham Linder
Superintendent

Jack R. Borsting
Provost

This research was partially supported by Grant AFORSR 74-2642 from the Air Force Office of Scientific Research and NSF Grant AG467 at the Naval Postgraduate School.

Reproduction of all or part of this report is authorized.

Prepared by:

Donald P. Gaver
DONALD P. GAVAR, Professor
Department of Operations Research

John P. Lehoczky
JOHN P. LEHOCZKY
Department of Statistics
Carnegie-Mellon University

Reviewed by:

Released by:

Michael G. Sovereign
MICHAEL G. SOVEREIGN, Chairman
Department of Operations Research

Robert R. Fossum
ROBERT R. FOSSUM
Dean of Research

| | | |
|---------------|--------------------------------|-------------------------------------|
| ACCESSION | White Section | <input checked="" type="checkbox"/> |
| NTIS | Blue Section | <input type="checkbox"/> |
| DOC | | <input type="checkbox"/> |
| UNANNOUNCED | | |
| JUSTIFICATION | | |
| BY | DISTRIBUTION AVAILABILITY CODE | |
| DATE | AVAIL. ON 1/14/80 | |

[Handwritten mark: A]

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--|---|---|
| 1. REPORT NUMBER | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| NPS55-77-16 | | |
| 4. TITLE (and Subtitle) | 5. TYPE OF REPORT & PERIOD COVERED | |
| A Diffusion Approximation Analysis of a General n-Compartment System. | Technical Report | |
| 6. AUTHOR(s) | 6. PERFORMING ORG. REPORT NUMBER | |
| Donald P. Gaver John P. Lehoczeky | | |
| 7. PERFORMING ORGANIZATION NAME AND ADDRESS | 8. CONTRACT OR GRANT NUMBER(s) | |
| Naval Postgraduate School Monterey, CA 93940 | Grant AFOSR 74-2642 NSF Grant AG467 | |
| 9. CONTROLLING OFFICE NAME AND ADDRESS | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS | |
| Naval Postgraduate School Monterey, CA 93940 | | |
| 11. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | 12. REPORT DATE | |
| | April 1977 | |
| | 13. NUMBER OF PAGES | |
| | 37 (12) 42 p. | |
| | 14. SECURITY CLASS. (of this report) | |
| | Unclassified | |
| | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE | |
| 16. DISTRIBUTION STATEMENT (of this Report) | | |
| Approved for public release; distribution unlimited | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) | | |
| Diffusion Theory, Pharmacokinetics, Probability, Compartments, Kinetics | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) | | |
| <p>A new approach to the stochastic analysis of general compartment models is presented. The analysis is based on the concept of diffusion approximations. The state of a compartment system is represented as the superposition of a deterministic process, characterized by a system of ordinary differential equations, and a random noise process characterized by stochastic differential equations. All transition rate parameters are permitted to be time dependent.</p> | | |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251 450 88

→ next
page

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

cont

Numerical solutions are presented for the two-compartment case, and compared with those of Cardenas and Matis (1975a). Extensions to non-linear compartment models are discussed.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

A DIFFUSION APPROXIMATION ANALYSIS
OF A GENERAL n -COMPARTMENT SYSTEM

by

John P. Lehoczky¹
Department of Statistics
Carnegie-Mellon University
Pittsburgh, Pennsylvania
15213

Donald P. Gaver²
Department of Operations Research
Naval Postgraduate School
Monterey, California
93940

ABSTRACT

A new approach to the stochastic analysis of general compartment models is presented. The analysis is based on the concept of diffusion approximations. The state of a compartment system is represented as the superposition of a deterministic process, characterized by a system of ordinary differential equations, and a random noise process characterized by stochastic differential equations. All transition rate parameters are permitted to be time dependent. Numerical solutions are presented for the two-compartment case, and compared with those of Cardenas and Matis (1975a). Extensions to non-linear compartment models are discussed.

¹Research supported in part by Grant AFOSR 74-2642 from the Air Force Office of Scientific Research.

²Research supported by NSF Grant AG467 at the Naval Postgraduate School.

1. Introduction

Quantitative representation of the distribution over time of a drug or pollutant in a human or animal body is now often carried out in terms of compartment models. Such elements as the blood, gut, liver, lean tissue, etc., are characterized as homogeneous compartments, within which the drug resides for a time, later to transit to another compartment, perhaps recycling but eventually vanishing because of evacuation or metabolism mechanisms.

Classical papers in this area, e.g. that by Bischoff et al. (1971) concerned with use of methotrexate in cancer therapy, proposed deterministic descriptions of flows between compartments, and stocks within, using differential equation systems. However, variations in drug concentrations over replicated experiments suggest a probabilistic or stochastic representation. Stochastic compartment analysis has undergone rapid development for several years, and might be categorized as follows. First, many papers have emphasized the mathematical formulation and exploration of the stochastic behavior of a variety of compartment models; papers of this type are exemplified by Bright (1973), Matis (1972, 1974, 1975a, 1975b), Marcus (1975), Matis and Hartley (1972), Purdue (1974a, 1974b, 1975), Rescigno (1973), Rubinow and Winzer (1971), Thakur, Rescigno, and Schafer (1972, 1973), and Thron (1972). Second, several authors have investigated problems of statistical inference for the transition rates in stochastic compartment models; see Burkinshaw and Marshall (1971), Cornfield, Steinfeld, and

Greenhouse (1970), Kodell and Matis (1976), Matis and Hartley (1972), Rodda et al. (1975), and Shah (1976). Last, the actual application of compartment models to biomedical, ecological, and chemical systems, as well as to pharmacokinetics, has been carried out, for example, by Metzler (1971), Rödenbeck et al. (1975), Sheppard (1962), Siegel, Cooper, and Meisner (1968) and Uppuluri and Bernard (1967). A large bibliography is given in Jacquez (1972).

The present paper falls into the first category: we suggest a new approach to stochastic compartment analysis based on mathematical diffusion processes; see Feller (1971) for an introduction, and Arnold (1973) and Gikhman and Skorohod (1971) for a systematic treatment. In short, we write stochastic differential equations to describe compartment concentration changes, and show that in a natural limit interesting joint distributions are jointly Gaussian or Normal. In the literature a variety of different compartment models have been formulated for one up to n -compartment, reversible or irreversible, open or closed, systems. These have generally been characterized by discrete-valued state variables for compartment contents, the latter changing according to multivariate birth-and-death processes. Kolmogorov forward equations for the transition probabilities are then derived, and solved by transforms.

Purdue (1974) notes the similarity of these models to those for infinite server queues. We develop and discuss this connection in the appendix of this paper. For such discrete models it is possible to write down expressions for the moments of compartment contents at

a fixed time; under certain conditions (Poisson inputs, transition rates depend linearly on compartment contents) the contents of the compartments are statistically independent and Poisson distributed. Cardenas and Matis (1975a,b) have extensively investigated such models. The approach taken here, that of replacing a discrete birth and death process with a continuous diffusion, is approximate. However, the results agree with the exact solution in the case of linear transition rates up to second moments. Furthermore, the limiting process may be shown rigorously to be normal, allowing the use of statistical inference procedures validated for normal distributions and processes. Perhaps more important is the fact that the diffusion approximation applies to situations with nonlinear transition Michaelis-Menten-type rates, where "exact" discrete-state methods yield cumbersome results; see McNeil and Schach (1973) for various examples concerning populations. It is reassuring to find that the diffusion approximations often agree exceptionally well with the birth and death solution; see Gaver and Lehoczký (1975,1976) for numerical comparisons.

2. A Model

We consider a general n -compartment model with time-dependent transition rates. Let $C_i(t)$, $i = 1, \dots, n$ represent the contents of compartments $1, \dots, n$ respectively at time t . We assume that the state of the system $\underline{C}(t) = (C_1(t), \dots, C_n(t))$ is a Markov process with transition probabilities given by

$$\begin{aligned}
 P(\Delta_i(t)=1, \Delta_j(t)=0, j \neq i | \underline{C}(t)) &= N\lambda_{0i}(t)dt + o(dt) \\
 &\quad i = 1, \dots, n \\
 P(\Delta_i(t)=-1, \Delta_j(t)=0, j \neq i | \underline{C}(t)) &= \lambda_{i0}(t) C_i(t)dt + o(dt) \\
 &\quad i = 1, \dots, n \\
 P(\Delta_i(t)=-1, \Delta_j(t)=+1, \Delta_k(t)=0, k \neq i, j | \underline{C}(t)) &= \lambda_{ij}(t) C_i(t)dt + o(dt) \\
 &\quad \text{for } 1 \leq i, j \leq n, i \neq j, \lambda_{ij} \geq 0.
 \end{aligned} \tag{1}$$

where $\Delta_i(t) = C_i(t + dt) - C_i(t)$, and N is a parameter that can be thought of as a dose level, eventually allowed to be large.

The first of the three transition probabilities represents an increase in compartment i of size 1 from the outside. The second represents a decrease in compartment i of size 1 to the outside. Finally, the third represents a transfer of size 1 from compartment i to compartment j .

The above formulation is rather general, and allows many different compartment systems to be studied simultaneously. It allows for an open system if $\lambda_{0i}(t) > 0$ for some $i = 1, \dots, n$ or a closed system if $\lambda_{0i}(t) = 0$ for $i = 1, m, n$. The flows can be either reversible if $\lambda_{ij}(t) > 0$ implies $\lambda_{ji}(t) > 0$ or irreversible if $\lambda_{ij}(t) > 0$ implies $\lambda_{ji}(t) = 0$.

We wish to study the case where $\sum_{i=1}^n C_i(t)$ is large, say proportional to some number N . A diffusion approximation arises when N becomes large and we expand $C_i(t)$ into terms of order N and \sqrt{N} . We can insure that $\sum_{i=1}^n C_i(t)$ is large in two different ways. For open systems we assume, as given in (1), that the transition probabilities into the system from the outside are themselves directly proportional to N . For closed systems where there is no input from the outside, we must assume that each compartment has contents proportional to N at time 0, that is $C_i(0) = Nc_i(0)$.

The effect of assuming $\sum_{i=1}^n C_i(t)$ to be proportional to N is that transitions of all types occur very frequently. As a result, in any short time interval there will be many transitions of each

type. Each type of transition represents a change of $+1$ or -1 in the contents of a certain compartment. In a short time interval the total change will be a sum of independent Bernoulli random variables, and, as such, normally distributed by virtue of the central limit theorem. Actually, much more is true. Focusing on the compartment process $\{C(t), t \geq 0\}$ over time rather than at a single fixed time we see that the process will have normally distributed increments and hence be Gaussian. This observation is the key to understanding the diffusion approximation approach. The mathematical development of this approach is not given here; see Kurtz (1971) and Barbour (1972), and the paper by Schach and McNeil (1973).

3. Mathematical Development

In this section we derive the diffusion approximation approach outlined above in a manner that seems intuitively appealing. The exposition will be in terms of (Ito-type) stochastic differential equations; although, as will appear subsequently, the stochastic differential equations describing stochastic variations in compartment contents are not ambiguous of interpretation. The derivation to be presented next is supplemented by further discussion, reserved for the Appendix, that relates the present theory to the approach by Barbour (1974), and also to infinite server queueing-type models.

Let, then, $dC_i(t) = C_i(t + dt) - C_i(t)$ represent the change in the contents of compartment i in time $(t, t + dt)$. The change $dC_i(t)$ may be viewed as a sum of independent random variables:

- (a) inputs from outside the system, with mean and variance $N\lambda_{0i}(t) dt$;
- (b) inputs from compartment j ($j \neq i$), with mean and variance $\lambda_{ji}(t) C_j(t) dt$;
- (c) outputs to compartment k ($k \neq i$), with mean and variance $\lambda_{ik}(t) C_i(t) dt$;
- (d) outputs from the system via compartment i , with mean and variance $\lambda_{i0}(t) C_i dt$.

Represent the change in contents of compartment i in the manner of diffusion, i.e. in terms of drift and diffusion terms:

$$\begin{aligned}
 dC_i(t) = & [N\lambda_{0i}(t) - \sum_{\substack{j=1 \\ j \neq i}}^n \lambda_{ij}(t) C_i(t) + \sum_{\substack{j=1 \\ j \neq i}}^n \lambda_{ji} C_j(t)] dt \\
 & + \sqrt{N\lambda_{0i}} dW_{0i}(t) - \sum_{\substack{j=0 \\ j \neq i}}^n \sqrt{\lambda_{ij}(t) C_i(t)} dW_{ij}(t) \\
 & + \sum_{\substack{j=1 \\ j \neq i}}^n \sqrt{\lambda_{ji}(t) C_j(t)} dW_{ji}(t)
 \end{aligned} \tag{2}$$

$i = 1, 2, \dots, n$; $C_i(0) = N\delta_{i0}$, and $\{W_{ij}(t), 0 \leq t, j \leq n, i \neq j\}$ is a family of standard Wiener processes, see Arnold (1974).

Expression (2) is motivated as follows: the drift term, in brackets, represents the expected change in $C_i(t)$ over (short) time interval $(t, t + dt)$, given the states of all compartments and the corresponding infinitesimal flow rates or transition probabilities. The remaining terms represent the various random components of change, modeled as Gaussian random variables; the latter is plausible as $N \rightarrow \infty$ on central limit theorem grounds. For instance, arrivals from outside the system in $(t, t + dt)$ are Poissonian, and hence for large N , nearly Gaussian, with mean and variance specified by (a) above, and this accounts for the first bracketed and unbracketed terms in (2).

It is shown in Kurtz (1971) that

$$\frac{C_i(t)}{N} \rightarrow c_i(t) \quad \text{as } N \rightarrow \infty \text{ in probability;} \quad (3)$$

see also Barbour (1974). By analogy with the central limit theorem, consider the normalized process

$$X_i(t) = \frac{C_i(t) - Nc_i(t)}{\sqrt{N}}; \quad (4)$$

hence express the concentration as

$$C_i(t) = Nc_i(t) + \sqrt{N} X_i(t). \quad (5)$$

In matrix form, $\underline{C}(t) = N\underline{c}(t) + \sqrt{N} \underline{X}(t)$.

The vector function $\tilde{N}_0(t)$ is referred to as the deterministic approximation, and $\sqrt{N} X(t)$ is a stochastic noise process superimposed upon the deterministic approximation. It remains to characterize $\tilde{c}(t)$ and $\tilde{X}(t)$. The stochastic process $\tilde{X}(t) = (X_1(t), \dots, X_n(t))$ defined by (4) satisfies a stochastic differential equation similar to (2), and it can be derived from (2) by an application of Ito's Lemma (Arnold, p. 90, or Gikhman and Skohorod, p. 24). The stochastic differential equation governing $X(t)$ is given by

$$d\tilde{X}(t) = \tilde{A}(t) \tilde{X}(t) dt + \tilde{B}(t) d\tilde{W}(t) - \sqrt{N} (\tilde{c}'(t) - \lambda_0(t) - \tilde{A}(t) \tilde{c}(t)) dt \quad (6)$$

where

$$\tilde{A}(t) = a_{ij}(t), \quad 1 \leq i, j \leq n$$

$$\text{with } a_{ij}(t) = \lambda_{ji}(t) \text{ for } i \neq j, a_{ii}(t) = - \sum_{\substack{j=0 \\ j \neq i}}^n \lambda_{ij}(t)$$

$$\lambda_0(t) = (\lambda_{01}(t), \dots, \lambda_{0n}(t))^T$$

$$\tilde{W}(t) = (W_{01}(t), \dots, W_{0n}(t), W_{10}(t), \dots, W_{n0}(t), W_{12}(t), \dots, W_{1n}(t), \dots, W_{n,n-1}(t))^T$$

$$\tilde{B}(t) = (b_{ik}(t)) \quad 1 \leq i \leq n, 1 \leq k \leq n(n+1)$$

with

$$b_{ii}(t) = \sqrt{\lambda_{0i}(t)} \quad i = 1, \dots, n$$

$$b_{i(n+i)}(t) = -\sqrt{\lambda_{i0}(t) C_i(t)/N}$$

$$\begin{aligned} b_{i(2n+(i-1)(n-1)+j-1)}(t) &= -b_{j(2n+(i-1)(n-1)+j-1)}(t) \\ &= -\sqrt{\lambda_{ij}(t) C_i(t)/N}, \quad j > i \end{aligned}$$

$$\begin{aligned} b_{i(2n+(i-1)(n-1)+j)}(t) &= -b_{j(2n+(i-1)(n-1)+j)}(t) \\ &= -\sqrt{\lambda_{ij}(t) C_i(t)/N}, \quad j < i \end{aligned}$$

and all other $b_{ik} = 0$.

We now let $N \rightarrow +\infty$ in (6). First, the terms in $B(t)$ involving $\sqrt{\lambda_{ij}(t) C_i(t)/N}$ converge to $\sqrt{\lambda_{ij}(t) c_i(t)}$. Second, equation (6) contains a term proportional to N . The coefficient of this term must be identically 0 for all $t \geq 0$ or (6) will not converge to a finite limit. Consequently $\zeta(t)$ must satisfy

$$\zeta'(t) = \lambda_0(t) + \underline{A}(t) \zeta(t) \quad (7)$$

and, given that $\zeta(t)$ satisfies (7), equation (6) becomes

$$d\underline{X}(t) = \underline{A}(t) \underline{X}(t) + \underline{B}(t) d\underline{W}(t) \quad (8)$$

where $C_i(t)/N$ terms are replaced by $c_i(t)$ in $\underline{B}(t)$. Note that the modelling ambiguity sometimes associated with stochastic differential equation representations, see Arnold (1973), Chap. 10, does not arise because the diffusion coefficient, $B(\cdot)$, of (8) does not involve \underline{X} .

The deterministic approximation to $\zeta(t)$ is given by $N\zeta(t)$ where $\zeta(t)$ is characterized by (7). This system of differential equations is stable in the sense of Liapunov because all eigenvalues of $A(t)$ have strictly negative real parts. The stochastic noise process superimposed on the deterministic approximation is given by $\sqrt{N}X(t)$ where $X(t)$ is characterized by (8). The stochastic differential equation given by (8) describes a nonstationary multivariate Ornstein-Uhlenbeck process. Much is known about such processes, see Arnold (1974); we will return to a discussion of the noise process in a later section.

4. The Deterministic Approximation

The system of differential equations given in (7) along with an initial configuration $\zeta(0)$ can be easily solved. If $\lambda_0(t) = 0$, the closed system case, then

$$\zeta(t) = \exp\left(\int_0^t A(s) ds\right) \cdot \zeta(0) \quad (9)$$

where $\exp(M) = I + M + M^2/2! + \dots$ for any square matrix M . For the open system, nonhomogeneous, case with $\lambda_0(t) \neq 0$ one first multiplies by $\exp(-\int_0^t A(s) ds)$ to obtain

$$\frac{d}{dt} \left(\exp\left(-\int_0^t A(s) ds\right) \cdot \zeta(t) \right) = \exp\left(-\int_0^t A(s) ds\right) \cdot \lambda_0(t) . \quad (10)$$

This equation can be integrated and the result multiplied by $\exp(\int_0^t \tilde{A}(s) ds)$ to give

$$\tilde{c}(t) = \int_0^t \exp(\int_u^t \tilde{A}(s) ds) \cdot \tilde{\lambda}_0(u) du + \exp(\int_0^t \tilde{A}(s) ds) \cdot \tilde{c}(0) \quad (11)$$

Equation (11) provides a complete solution to the problem of describing $\tilde{c}(t)$. Simplifications are not possible in general; however, equation (11) can always be solved numerically by standard numerical integration techniques. We can consider a special case which allows equation (11) to be substantially simplified. Suppose we assume that all transition probabilities, $\lambda_{ij}(t)$, $1 \leq i \leq n$, $0 \leq j \leq n$, $i \neq j$, exhibit the same time dependence, or that $\lambda_{ij}(t) = \lambda_{ij} f(t)$ where $f(t)$ is some function of time. The matrix $\tilde{A}(t)$ now has the form

$$\tilde{A}(t) = \tilde{A} f(t) \quad (12)$$

where

$$\tilde{A} = (a_{ij}) \quad \begin{cases} a_{ij} = \lambda_{ji} & j \neq i \\ a_{ii} = - \sum_{\substack{j=0 \\ j \neq i}}^n \lambda_{ij} \end{cases}$$

Assume that \tilde{A} can be diagonalized and has eigenvalues $\theta_1, \theta_2, \dots, \theta_n$, left eigenvectors $\tilde{\lambda}_1, \dots, \tilde{\lambda}_n$, and right eigenvectors $\tilde{x}_1, \dots, \tilde{x}_n$. This gives

$$\underline{\dot{A}}(t) = \underline{R}\underline{D}(t)\underline{L} \quad \text{with} \quad \underline{L}\underline{R} = \underline{I} \quad (13)$$

where $\underline{R} = (r_1, r_2, \dots, r_n)$, $\underline{L} = (l_1, \dots, l_n)'$, and $\underline{D}(t)$ is a diagonal matrix with i th entry $\theta_i f(t)$. Equation (13) can be integrated to give

$$\int_u^t \underline{\dot{A}}(s) ds = \underline{R}\underline{E}(u, t)\underline{L} \quad (14)$$

with $\underline{E}(u, t)$ a diagonal matrix having elements $(\theta_i \int_u^t f(s) ds)$. Finally (14) can be exponentiated to give

$$\exp \int_u^t \underline{\dot{A}}(s) ds = \underline{R}\underline{G}(u, t)\underline{L} \quad (15)$$

where $\underline{G}(u, t)$ is a diagonal matrix with the i th diagonal element $\exp(\theta_i \int_u^t f(s) ds)$. Equation (15) can be plugged back into (11) to give

$$\underline{C}(t) = \underline{R} \int_0^t \underline{G}(u, t) \cdot \underline{L} \cdot \underline{\lambda}_0(u) du + \underline{R}\underline{G}(0, t)\underline{L} \cdot \underline{C}(0) \quad (16)$$

We illustrate the use of (16) by considering an example discussed by Cardenas and Matis (1975a). In that paper the authors consider a case with time-dependent transitions for a closed two-compartment system. For closed systems $\underline{\lambda}_0(u) = 0$ and only the second term of (16) need be calculated. In this case it is assumed that

$$\lambda_{01}(t) = \lambda_{02}(t) = 0 ,$$

$$\lambda_{12}(t) = .3/(t+1) ,$$

$$\lambda_{10}(t) = .5/(t+1) ,$$

$$\lambda_{21}(t) = .4/(t+1) ,$$

and

$$\lambda_{20}(t) = .6/(t+1) .$$

$$A(s) = \begin{pmatrix} -.8 & .4 & \frac{1}{s+1} \\ .3 & -1 & \end{pmatrix}$$

and

$$A = \begin{pmatrix} -.8 & .4 \\ .3 & -1 \end{pmatrix} , \quad f(s) = \frac{1}{s+1} .$$

The matrix A has eigenvalues $\theta_1 = -.9 + \sqrt{.13}$, $\theta_2 = -.9 - \sqrt{.13}$. Also $\exp(\theta_i \int_0^t f(s) ds) = (1+t)^{\theta_i}$. It is easy to calculate

$$\xi(t) = \begin{pmatrix} 1 & 1 \\ a/4 & b/4 \end{pmatrix} \begin{pmatrix} (1+t)^{\theta_1} & 0 \\ 0 & (1+t)^{\theta_2} \end{pmatrix} \begin{pmatrix} .12/(.12+a^2) & .4a/(.12+a^2) \\ .12/(.12+b^2) & .4b/(.12+b^2) \end{pmatrix} \xi(0)$$

$$a = -.1 + \sqrt{.13} \tag{17}$$

$$b = -.1 - \sqrt{.13}$$

Equation (17) may be expanded to yield

$$\begin{aligned}
c_1(t) = & (.6386750491c_1(0) + .5547001962c_2(0)) (1+t)^{\theta_1} \\
& + (.361324909c_1(0) - .5547001962c_2(0)) (1+t)^{\theta_2}
\end{aligned}
\tag{18}$$

$$\begin{aligned}
c_2(t) = & (.4160251472c_1(0) + .3613249509c_2(0)) (1+t)^{\theta_1} \\
& + (-.4160251472c_1(0) + .6386750491c_2(0)) (1+t)^{\theta_2}
\end{aligned}$$

with

$$\theta_1 = - .5394448725$$

$$\theta_2 = -1.2605551275$$

Transition probabilities for this model were approximated by Cardenas and Matis using a truncated infinite series approach. These transition probabilities can be derived directly from (18) by establishing appropriate initial conditions. For example, by setting $c_1(0) = 1$, $c_2(0) = 0$, $c_1(t)$ and $c_2(t)$ become $p_{11}(t)$ and $p_{12}(t)$, respectively. By setting $c_1(0) = 0$, $c_2(0) = 1$, $c_1(t)$ and $c_2(t)$ become $p_{21}(t)$ and $p_{22}(t)$, respectively. The results derived from (18) are compared with the approximations of Cardenas and Matis in Table 1. These results illustrate the accuracy of the deterministic approximations in (18), since the two answers agree to at least 8 significant figures. Furthermore, the numbers derived from (18) always exceed the Cardenas and Matis results. This is reasonable since the latter approximation is derived by truncation and is hence an underestimate of the true value.

The results in (18) illustrate another very important point. Compartment modelling has been criticized because it only allows levels which are mixtures of exponentials in t when, in fact, these models often do not fit data well (See Marcus (1975) p. 339 and Matis (1972)). Much more complicated functions are required to give adequate fits including gamma, Pareto, Pareto-exponential, and first passage density functions. One can see that all of these types of functions can be produced by appropriate choice of $f(t)$. Mixtures of exponentials arise when $f(t) \equiv 1$; however, an enormous class of level functions can be produced by varying choices of $f(t)$. The introduction of time dependent transitions not only adds realism but produces models which can fit real data sets.

| | t | Equation (18) | Cardenas and Matis 4-term approximation |
|-------------|---|---------------|--|
| $P_{11}(t)$ | 1 | .5902421829 | .5902421826 |
| | 2 | .4435620915 | .4435620872 |
| | 3 | .3652902915 | .3652902729 |
| | 4 | .3155676392 | .3155675836 |
| | 5 | .2807030964 | .2807029846 |
| $P_{22}(t)$ | 1 | .5151767471 | .5151767467 |
| | 2 | .3596617846 | .3596617805 |
| | 3 | .2823115397 | .2823115195 |
| | 4 | .2356325933 | .2356325396 |
| | 5 | .2041834275 | .2041833201 |
| $P_{21}(t)$ | 1 | .1501308716 | .1501308715 |
| | 2 | .1678006137 | .1678006133 |
| | 3 | .1659575075 | .1659575068 |
| | 4 | .1598700918 | .1598700879 |
| | 5 | .1530393378 | .1530393292 |
| $P_{12}(t)$ | 1 | .1125981537 | .1125981539 |
| | 2 | .1258504603 | .1258504600 |
| | 3 | .1244681306 | .1244681301 |
| | 4 | .1199025689 | .1199025659 |
| | 5 | .1147795033 | .1147794969 |

TABLE 1: A Comparison of the Deterministic Approximation (18) with the Cardenas and Matis Four Term Approximation.

5. The Noise Process

The stochastic elements of the compartment process $\{\zeta(t), t \geq 0\}$ are embodied in the nonstationary multivariate Ornstein-Uhlenbeck process described by (8). We summarize a few key results about such processes. These results are well known and available in Arnold, Section 8.2. First, we assume $\tilde{x}(0) = \underline{0}$ that is the initial condition is nonstochastic and embodied in $\zeta(0)$. The marginal distribution of $\tilde{x}(t)$ is

$$\tilde{x}(t) \sim \mathcal{N}_n(\underline{0}, \Sigma(t)) \quad (19)$$

that is an n -dimensional multivariate normal distribution where $\Sigma(t)$ is the unique nonnegative definite solution of the matrix Riccati equation

$$\Sigma'(t) = \underline{A}(t) \Sigma(t) + \Sigma(t) \underline{A}^T(t) + \underline{B}(t) \underline{B}^T(t) \quad (20)$$

Since $\zeta(t) = N\zeta(t) + \sqrt{N} \tilde{x}(t)$ we find

$$\zeta(t) \sim \mathcal{N}_n(N\zeta(t), N\Sigma(t)) \quad (21)$$

Let $\underline{B}(t) \underline{B}^T(t) = \underline{D}(t)$ and $\underline{D}(t) = (d_{ij}(t))$ with

$$\begin{aligned} d_{ii}(t) &= (\lambda_{0i}(t) + \sum_{j=0}^n \lambda_{ij}(t) c_i(t) + \sum_{j=1}^n \lambda_{ji}(t) c_j(t)) \\ d_{ij}(t) &= -(\lambda_{ij}(t) c_i(t) + \lambda_{ji}(t) c_j(t)) \end{aligned} \quad (22)$$

The covariance matrix $\underline{\Sigma}(t)$ can be calculated by various numerical methods applied to (20). We illustrate the calculation in the case $n = 2$ studied by Cardenas and Matis.

$$\underline{A}(t) = \begin{pmatrix} -(\lambda_{10}(t) + \lambda_{12}(t)) & \lambda_{21}(t) \\ \lambda_{12}(t) & -(\lambda_{20}(t) + \lambda_{21}(t)) \end{pmatrix}$$

$$\underline{\Sigma}(t) = \begin{pmatrix} \sigma_{11}(t) & \sigma_{12}(t) \\ \sigma_{12}(t) & \sigma_{22}(t) \end{pmatrix}$$

Equation (20) becomes

$$\dot{\underline{s}}(t) = \underline{G}(t) \underline{s}(t) + \underline{H}(t) \quad (23)$$

where

$$\underline{s}(t) = (\sigma_{11}(t), \sigma_{12}(t), \sigma_{22}(t))'$$

$$\underline{G}(t)$$

$$= \begin{pmatrix} -2(\lambda_{10}(t) + \lambda_{12}(t)) & 2\lambda_{21}(t) & 0 \\ \lambda_{12}(t) & -(\lambda_{10}(t) + \lambda_{20}(t) + \lambda_{12}(t) + \lambda_{21}(t)) & \lambda_{21}(t) \\ 0 & 2\lambda_{12}(t) & -2(\lambda_{20}(t) + \lambda_{21}(t)) \end{pmatrix}$$

$$\underline{H}(t) = \begin{pmatrix} \lambda_{01}(t) + (\lambda_{10}(t) + \lambda_{12}(t)c_1(t) + \lambda_{21}(t)c_2(t)) \\ -(\lambda_{12}(t)c_1(t) + \lambda_{21}(t)c_2(t)) \\ \lambda_{02}(t) + \lambda_{12}(t)c_1(t) + (\lambda_{20}(t) + \lambda_{21}(t)c_2(t)) \end{pmatrix}.$$

Equation (23) can be solved in exactly the same manner as (7) with initial conditions $\underline{s}(0) = \underline{0}$, that is $\underline{C}(0)$ is deterministic. We find

$$\underline{s}(t) = \int_0^t \exp\left(\int_u^t \underline{G}(s) ds\right) \cdot \underline{H}(u) du . \quad (24)$$

For the case considered by Cardenas and Matis

$$\underline{G}(t) = \frac{1}{1+t} \begin{pmatrix} -1.6 & .8 & 0 \\ .3 & -1.8 & .4 \\ 0 & .6 & -2 \end{pmatrix} + \frac{1}{1+t} \underline{G} .$$

The matrix \underline{G} has eigenvalues ϕ_1, ϕ_2, ϕ_3 where

$$\phi_1 = 2\theta_1 = -1.978889745$$

$$\phi_2 = \theta_1 + \theta_2 = -1.8,$$

$$\phi_3 = 2\theta_2 = -2.521110255$$

$$\underline{H}(u) = \begin{pmatrix} k_{11}(u+1)^{\theta_1-1} + k_{12}(u+1)^{\theta_2-1} \\ k_{21}(u+1)^{\theta_1-1} + k_{22}(u+1)^{\theta_2-1} \\ k_{31}(u+1)^{\theta_1-1} + k_{32}(u+1)^{\theta_2-1} \end{pmatrix}$$

where the k_{ij} coefficients are computing from (18). The other factor in the integrand is given by

$$\exp \int_u^t \underline{G}(s) ds = \underline{R} \begin{pmatrix} \left(\frac{1+t}{1+u}\right)^{\phi_1} & 0 & 0 \\ 0 & \left(\frac{1+t}{1+u}\right)^{\phi_2} & 0 \\ 0 & 0 & \left(\frac{1+t}{1+u}\right)^{\phi_3} \end{pmatrix} \underline{L}$$

with

$$R = \begin{pmatrix} 1 & 1 & 1 \\ .6513878188 & -.25 & -1.151387819 \\ .4243060905 & -.75 & 1.32569391 \end{pmatrix}$$

$$L = \begin{pmatrix} .407957184 & .7085463502 & .3076923077 \\ .4615384615 & -.3076923077 & -.6153846153 \\ .1305557202 & -.4998540425 & 1 \end{pmatrix}$$

$\underline{s}(t)$ can now be easily found. Each of the three variance-covariance terms is a linear combination of terms of the form

$$(1+t)^{\phi_i} ((1+t)^{\phi_j - \phi_i} - 1) / (\phi_j - \phi_i) .$$

The Ornstein Uhlenbeck process described by (8) also has a known covariance function $k(s,t) = \text{Cov}(\underline{X}(s), \underline{X}(t))$. The function $k(s,t)$ is given by the equation (T denotes transpose)

$$k(s,t) = \underline{\Phi}(s) \int_0^{\min(s,t)} \underline{\Phi}^{-1}(u) \underline{B}(u) \underline{B}^T(u) (\underline{\Phi}^{-1}(u))^T du \underline{\Phi}^T(t) \quad (25)$$

with

$$\underline{\Phi}(u) = \exp\left(\int_0^u A(s) ds\right) .$$

We do not use the covariance function in this paper; however, this function is important in the development of a statistical analysis of data from a compartment model. One may have data, say readings of drug concentrations in the blood stream, and wish to

estimate the parameters of the model, the λ_{ij} 's. Given the data have a multivariate normal distribution, the likelihood function can be written and estimators derived. The papers of Hartley and Matis (1971) and Kodell and Matis (1976) provide analyses in special cases. The results of this paper indicate that these kinds of results can be carried much further into far more general compartment models.

6. Steady State

In the case $\lambda_0(t) \neq 0$, an open system, the system will approach steady state if $\lambda_{ij}(t) \rightarrow \lambda_{ij}$ as $t \rightarrow +\infty$. Steady state is a condition whereby the probability distribution which describes the marginal distribution of the system becomes time homogeneous even though the actual contents of the compartments continue to vary according to (8). The steady-state solution for a simple example is described in the Appendix.

The process $\{\tilde{C}(t), t \geq 0\}$ in steady state can also be approximated by $\tilde{N}\tilde{C} + \sqrt{\tilde{N}}\tilde{X}(t)$ where

$$0 = \tilde{\lambda}_0 + \tilde{A}\tilde{C} \quad \text{or} \quad \tilde{C} = -\tilde{A}^{-1}\tilde{\lambda}_0 \quad (26)$$

and $\{\tilde{X}(t), t \geq 0\}$ satisfies

$$d\tilde{X}(t) = \tilde{A}\tilde{X}(t) + \tilde{B} d\tilde{W}(t) \quad (27)$$

a stationary multivariate Ornstein-Uhlenbeck process. This means that

$$\tilde{C}(t) \sim \mathcal{N}_n(-\tilde{N}\tilde{A}^{-1}\tilde{\lambda}_0, \tilde{N}\tilde{\Sigma}) \quad (28)$$

where $\tilde{\Sigma}$ is the unique nonnegative definite solution of

$$\tilde{A}\tilde{\Sigma} + \tilde{\Sigma}\tilde{A}^T = -\tilde{B}\tilde{B}^T. \quad (29)$$

As an illustration we consider a two compartment open system, either reversible or irreversible

$$\begin{aligned} \tilde{A} &= \begin{pmatrix} -(\lambda_{10} + \lambda_{12}) & \lambda_{21} \\ 0 & -(\lambda_{20} + \lambda_{21}) \end{pmatrix} \\ \tilde{B}\tilde{B}^T &= \begin{pmatrix} \lambda_{01} + (\lambda_{10} + \lambda_{12})c_1 + \lambda_{21}c_2 & \\ -(\lambda_{12}c_1 + \lambda_{21}c_2) & \\ \lambda_{02} + \lambda_{12}c_1 + (\lambda_{20} + \lambda_{21})c_2 & \end{pmatrix} \\ c &= \frac{\begin{pmatrix} \lambda_{20} + \lambda_{21} & \lambda_{12} \\ \lambda_{21} & \lambda_{10} + \lambda_{12} \end{pmatrix} \begin{pmatrix} \lambda_{01} \\ \lambda_{02} \end{pmatrix}}{(\lambda_{20} + \lambda_{21})(\lambda_{10} + \lambda_{12}) - \lambda_{12}\lambda_{21}} \end{aligned} \quad (30)$$

$$\begin{pmatrix} \sigma_{11} \\ \sigma_{12} \\ \sigma_{22} \end{pmatrix} = \begin{pmatrix} 2(\lambda_{10} + \lambda_{12}) & 2\lambda_{21} & 0 \\ \lambda_{12} & (\lambda_{10} + \lambda_{20} + \lambda_{12} + \lambda_{21}) & -\lambda_{21} \\ 0 & -2\lambda_{12} & 2(\lambda_{20} + \lambda_{21}) \end{pmatrix}^{-1} \quad \text{BB}' \quad (31)$$

Equations (30) and (31) thus complete the description of the multivariate normal density given in (28). While the $n = 2$ case can be done in closed form, the general case can be easily solved by numerical methods.

7. Extension to Nonlinear Models

The technique of diffusion approximations outlined and applied to general linear compartment models in the previous section also provides a tractable approach to handling nonlinear models. The compartment modelling literature is vast and the area is still in rapid development. Nevertheless, papers involving a stochastic process approach assume a linear system, that is, one in which transition rates from i to j are proportional to the contents of the i th compartment. It is unlikely that real pharmacokinetic processes operate so simply. Rather, the transition rates are more likely to be complicated functions of the contents of several compartment levels, and of time. Such complicated models have not appeared in the literature, perhaps because they present serious mathematical intractabilities using the standard Kolmogorov forward equations analysis.

The method of diffusion approximations can effectively handle nonlinear transition rates. The deterministic approximation will be a system of nonlinear ordinary differential equations. This will only rarely be solvable in closed form. Nevertheless, the numerical solution of such systems is a standard topic in numerical analysis. The important observation is that the stochastic differential equation representing the noise process superimposed upon the deterministic process will be nonstationary but linear (often Ornstein-Uhlenbeck). This means that the diffusion approximation approach will yield analytic results for nonlinear systems, since many results are readily available for linear stochastic differential equations (see Arnold, Chapter 8). It is hoped that this approach will encourage pharmacokinetic researchers to consider introducing nonlinear models should an analysis of data indicate the need.

Two additional points should be made. First, the analysis presented can be easily carried out when the process is represented in terms of volume and concentration rather than in terms of total contents. Second, the method of diffusion approximations illustrates that the compartment system contents will have a marginal Gaussian distribution. As such, one is in a position to give a statistical analysis of such a system either to estimate transition rates or, indeed, to design an experiment to make proper inferences about such parameters.

REFERENCES

- Arnold, L., Stochastic Differential Equations: Theory and Applications, Wiley, New York, 1973.
- Atkins, G. L. Multicompartment Models for Biological Systems, Methuen, London, 1969.
- Barbour, A. D., "On a Functional Central Limit Theorem for Markov Population Processes," Adv. Appl. Prob., Vol. 6 pp. 21-39, 1974.
- Bartlett, M. S., "Some evolutionary stochastic processes," J. R. Stat. Soc., B, Vol. 11, pp. 211-229, 1949.
- Bernard, S. R., Shenton, L. R., and Uppuluri, V. R. R., "Stochastic Models for the Distribution of Radioactive Material in a Corrected System of Compartments," in Proc. 5th Berkeley Symp. Math. Stat. Prob., Vol. IV, pp. 481-510, 1965.
- Bischoff, K. D., Dedrick, R. L., Zaharko, D. S., et al., "Methotrexate Pharmacokinetics," J. Pharm. Sci., Vol. 60, pp. 1128-1133, 1971.
- Bright, P. B., "The Volumes of Some Compartment Systems With Sampling and Loss From One Compartment," Bull. Math. Biol., Vol. 35, pp. 69-79, 1973.
- Burkinshaw, L. and Marshall, D. H., "Accuracy of Estimation of the Parameters in a Model of Calcium Tracer Kinetics," J. Theor. Biol., Vol. 30, pp. 255-265, 1971.
- Cardenas, M. and Matis, J. H., "On the Stochastic Theory of Compartments: Solution for N-Compartment Systems With Irreversible, Time-Dependent Transition Probabilities," Bull. Math. Biol., Vol. 36, pp. 480-504, 1974.
- Cardenas, M. and Matis, J. H., "On the Time-Dependent Reversible Stochastic Compartmental Model I: The General Two-Compartment System," Bull. Math. Biol., Vol. 37, pp. 505-519, 1975a.
- Cardenas, M. and Matis, J. H., "On the Time-Dependent Reversible Stochastic Compartmental Model: II. A Class of N-Compartment Systems," Bull. Math. Biol., Vol. 37, 1975b.
- Cornfield, J., Steinfeld, J., and Greenhouse, S. W., "Models for the Interpretation of Experiments Using Tracer Compounds," Biometrics, Vol. 16, pp. 212-234, (1960).
- Feller, W., An Introduction to Probability Theory and Its Application, John Wiley and Sons, New York, Vol. I, 1959, Vol. II, 1971.

References (cont'd)

- Gaver, D. P. and Lehoczky, J. P., Gaussian Approximation to Service Problems: A Communication System Example. J. Appl. Prob. to appear.
- Gaver, D. P. and Lehoczky, J. P., "A Diffusion Approximation for a Repairman Problem with Two Types of Repair," to appear Management Science.
- Gikhman, I. I., and Skorokhod, A. V., Stochastic Differential Equations, Springer-Verlag, New York, 1971.
- Hearon, J. F., "Resident Times in Compartmental Systems and the Moments of a Certain Distribution," Math. Biosciences, Vol. 15, pp. 69-77, 1972.
- Jacquez, J. A., Compartmental Analysis in Biology and Medicine, Elsevier Publishing Co., Amsterdam, 1972.
- Kingman, J. F. C., "Markov population processes." J. Appl. Prob., Vol. 6, No. 1, pp. 1-18, 1969.
- Kodell, R. L. and Matis, J. H., "Estimating the Rate Constants in a Two-Compartment Stochastic Model," to appear Biometrics, 1976.
- Kurtz, T. G., "Limit Theorems for Sequences of Jump Markov Processes Approximating Ordinary Differential Processes," J. Appl. Prob., Vol. 8, pp. 344-356, 1971.
- Marcus, A. H., "Power Laws in Compartmental Analysis, Part I: A Unified Stochastic Model," Math Biosciences, Vol. 23, pp. 337-350, 1975.
- Matis, J. H., "Gamma Time Dependency in Blaxter's Compartmental Model," Biometrics, Vol. 28, pp. 597-602, 1972.
- Matis, J. H., and Hartley, H. O., "Stochastic Compartmental Analysis: Model and Least Squares Estimation from Time Series Data," Biometrics, Vol. 28, pp. 77-97, 1972.
- McNeil, D. R., and Schach, S., "Central Limit Analogues for Markov Population Processes," J. Royal Stat. Soc. (B), Vol. 35, No. 1, 1973.
- Metzler, C. M., "Usefulness of the Two-Compartment Open Model in Pharmacokinetics," Journal of American Statistical Association, Vol. 66, pp. 49-53, 1971.

References (cont'd)

- Purdue, P., "Stochastic Theory of Compartments," Bull. Math. Biology, Vol. 36, pp. 305-309, 1974a.
- Purdue, P., "Stochastic Theory of Compartments: One and Two Compartment Systems," Bull. Math. Biol., Vol. 36, pp. 577-587, 1974b.
- Purdue, P., "Stochastic Theory of Compartments: An Open, Two-Compartment, Reversible System with Independent Poisson Arrivals," Bull. Math. Biol., Vol. 37, pp. 269-275, 1975.
- Rescigno, A., "Transfer Times in Tracer Experiments," J. Theor. Biol., Vol. 39, pp. 9-27, 1973.
- Rodda, B. E., Sampson, C. B., and Smith, D. W., "The One-Compartment Open Model: Some Statistical Aspects of Parameter Estimation," Applied Statistics, Vol. 24, pp. 309-318, 1975.
- Rödenbeck, M., Ziegler, A., Marek, H., and Friedrich, T.H., "Computer Analysis of a Tracer-kinetic Model of the Collagenic Metabolism of the Rat's Aorta," J. Theor. Biol., Vol. 49, pp. 191-200, 1975.
- Rubinow, S. I. and Winzer, A., "Compartment Analysis: An Inverse Problem," Math. Biosciences, Vol. II, pp. 203-247, 1971.
- Shah, B. K., "Data Analysis Problems in the Area of Pharmacokinetics Research," Biometrics, Vol. 32, pp. 145-157, 1976.
- Sheppard, C. W., Basic Principles of the Tracer Method, Wiley, New York, 1962.
- Siegel, C., Cooper, T. B., and Meisner, M., "Mathematical Models of the Thyroid: Deterministic and Stochastic," in Computers and Electronic Devices in Psychiatry, (N. Kline and E. Laska, eds.) Grune and Stratton, New York, pp. 100-108, 1968.
- Thakur, A. K., Rescigno, A., and Schafer, D., "On the Stochastic Theory of Compartments, I. A Single Compartment System," Bull. Math. Biol., Vol. 34, pp. 53-63, 1972.
- Thakur, A. K., Rescigno, A., and Schafer, D., "On the Stochastic Theory of Compartments, II. Multi-compartment Systems," Bull. Math. Biol., Vol. 35, pp. 263-271, 1973.

References (cont'd)

- Thron, C. L., "Structure and Kinetic Behavior of Linear Multicompartment Systems," Bull. Math. Biophys., Vol. 34, pp. 277-291, 1972.
- Uppuluri, V. R. R. and Bernard, S. R., "A Two-Compartment Model With a Random Connection," In Compartments, Pools and Spaces in Medical Physiology (P. E. E. Bergner and C. C. Lushbaugh, eds.) Brookhaven Symp. Series, No. 11, U.S.A.E.C., 1967.

APPENDIX

The purpose of this appendix is to further justify the earlier analysis, in particular, equations (7) and (8) and their implications. We do this in certain tractable simple cases, namely, those of one and two components.

1. The Theory of Kurtz (1971) and of Barbour (1974).

In Section 2 of Barbour (1974), hereafter called B, assumptions are given that imply convergence of the sequence of processes

$$\tilde{y}_N(t) = \sqrt{N} \left(\frac{\tilde{C}_N(t)}{N} - \tilde{c}(t) \right) = \frac{\tilde{C}_N(t) - N\tilde{c}(t)}{\sqrt{N}}, \quad N = 1, 2, 3, \dots, \quad (A-1)$$

to the diffusion $\tilde{y}(t)$, as $N \rightarrow \infty$ in $D(0, T)$, i.e. the space of all right-continuous functions with left-hand limits. Note that the $\tilde{y}(t)$ of B will turn out to be the same as our stochastic noise process $\tilde{x}(t)$. We shall verify these assumptions for illustrative cases.

First consider the sequence of continuous parameter Markov processes $\tilde{x}_N(t) = \tilde{C}_N(t)/N$ on $[0, T]$, where $\tilde{C}_N(t)$ is the many-compartment model whose transitions are described by our (1). Notice that the changes in $\tilde{x}_N(t)$ are of size $0, \pm 1/N$, and that the infinitesimal transition rates are of the form $N\lambda_{0i}$, and $N\lambda_{ij}(C_i(t)/N)$, as specified by (2.1, B). We now discuss the assumption and the Theorem K of B for these illustrative situations.

Single-Compartment Model. Let $\lambda_{01}(t) = \lambda_{01}$, and $\lambda_{10}(t) = \lambda_{10}$ constant and positive, and $n = 1$, which represents a single-compartment system.

It is easily seen that there is a unique function $\xi(t)$ satisfying, for $0 \leq t \leq T$, the natural deterministic equation, (2.2,B):

$$\xi'(t) = \lambda_{01} - \lambda_{10}\xi(t). \quad (A-2)$$

The solution is, putting $\xi(0) = c(0)$,

$$\xi(t) = c(0)e^{-\lambda_{10}t} + \frac{\lambda_{01}}{\lambda_{10}}(1 - e^{-\lambda_{10}t}) = \left(c(0) - \frac{\lambda_{01}}{\lambda_{10}}\right)e^{-\lambda_{10}t} + \frac{\lambda_{01}}{\lambda_{10}}, \quad (A-3)$$

which is unique; hence B's Assumption A is justified. Suppose $0 < c(0) < \lambda_{01}/\lambda_{10}$ for example; other cases may be treated in analogous fashion. Then, in the terminology of Barbour (1974),

$$S = \left[c(0), c(0)e^{-\lambda_{10}T} + \frac{\lambda_{01}}{\lambda_{10}}(1 - e^{-\lambda_{10}T}) \right] \quad (A-4)$$

and

$$S^\varepsilon = \left[c(0) - \varepsilon, c(0)e^{-\lambda_{10}T} + \frac{\lambda_{01}}{\lambda_{10}}(1 - e^{-\lambda_{10}T}) + \varepsilon \right], \quad (A-5)$$

where $0 < \varepsilon < c(0)$ suffices. Clearly the transition rates are multinomial within S^ε , and B's Assumption B is also justified. Now Theorem K follows, and implies that $\underline{y}_N(t) = \sqrt{N}(\underline{x}_N(t) - \xi(t))$ converges to the diffusion $\underline{y}(t)$, $\underline{y}(0) = 0$, with the characteristic

function $\phi(\theta, t) = E[e^{i\theta y(t)}]$ satisfying the differential equation

$$\frac{\partial \phi}{\partial t} = -\theta \lambda_{10} \frac{\partial \phi}{\partial \theta} - \frac{1}{2} \theta^2 [\lambda_{01} + \lambda_{10} \xi(t)] \phi \quad (A-6)$$

recognizable as describing the Ornstein-Uhlenbeck process whose stochastic differential equation is

$$dy = -\lambda_{10} y dt + \sqrt{\lambda_{01} + \lambda_{10} \xi(t)} dw, \quad (A-7)$$

this being the same s.d.e. as that obeyed by our noise process X for the present setup. Hence the procedure leading to our equations (7) and (8) yields the same results as those rigorously established by Kurtz, adapted by Barbour.

Note, too, that a steady-state solution will exist: simply set $\xi' = 0$ in (A-2) to discover the deterministic component: $\xi(\infty) = \lambda_{01}/\lambda_{10}$. The stochastic noise has, from (A-6) or (A-7), variance equal to the mean, namely, $\lambda_{01}/\lambda_{10}$. Thus the steady-state compartment content is, according to our theory, approximately $\eta(N\lambda_{01}/\lambda_{10}, N\lambda_{01}/\lambda_{10})$, as N becomes large. This is in accord with the fact that the exact distribution is Poisson $(N\lambda_{01}/\lambda_{10})$, as is well-known, and shown again subsequently in this Appendix.

Two-Component Model. Here $\underline{C}(t) = (C_1(t), C_2(t))$, the allowed transitions and rates are summarized below.

| <u>Transitions</u> | <u>Transition Rates</u> |
|------------------------------|---|
| $C_1(t), C_2(t) \rightarrow$ | |
| a) $C_1(t) + 1, C_2(t)$ | $N\lambda_{01}$ |
| b) $C_1(t) - 1, C_2(t)$ | $N\lambda_{10} \left(\frac{C_1(t)}{N} \right)$ |
| c) $C_1(t) - 1, C_2(t) + 1$ | $N\lambda_{12} \left(\frac{C_1(t)}{N} \right)$ |
| d) $C_1(t), C_2(t) - 1$ | $N\lambda_{20} \left(\frac{C_2(t)}{N} \right)$ |

(A-8)

State scaling, as in (2.1,B) alters the ± 1 changes to $\pm 1/N$.

The function $\underline{\xi}(t) = (\xi_1(t), \xi_2(t))$ satisfies the differential equations

$$\xi_1' = \lambda_{01} - (\lambda_{10} + \lambda_{12})\xi_1$$

(A-9)

$$\xi_2' = \lambda_{12}\xi_1 - \lambda_{20}\xi_2$$

For simplicity, we have not allowed entries into compartment 2 from outside, and have also omitted back flow from compartment 2 to compartment 1. The two equations are readily solved to produce a unique solution for all $t \geq 0$:

$$\xi_1(t) = c_1(0) \exp[-(\lambda_{10} + \lambda_{12})t] + \frac{\lambda_{01}}{\lambda_{10} + \lambda_{12}} [1 - \exp(-(\lambda_{10} + \lambda_{12})t)]$$

$$\begin{aligned}
\xi_2(t) = & c_{20} \exp(-\lambda_{20}t) + \frac{\lambda_{01}}{\lambda_{10} + \lambda_{12}} \frac{\lambda_{12}}{\lambda_{20}} [1 - \exp(-\lambda_{20}t)] \\
& + \left[c_1(0) - \frac{\lambda_{01}}{\lambda_{10} + \lambda_{12}} \right] \frac{\lambda_{12}}{\lambda_{20} - (\lambda_{10} + \lambda_{12})} \\
& \times [\exp(-(\lambda_{10} + \lambda_{12})t) - \exp(-\lambda_{20}t)] \quad (A-10)
\end{aligned}$$

Thus Assumption A is satisfied, with $c_1(0), c_2(0) > 0$. Assumption B is also satisfied with

$$0 < \varepsilon \leq \min[\underline{\xi}_1(T), \underline{\xi}_2(T)] \quad (A-11)$$

where

$$\bar{\xi}_i(T) = \min_{0 \leq t \leq T} \xi_i(t) > 0, \quad i = 1, 2;$$

It is clear from the differential equations that such values will always exist. Hence

$$\tilde{y}_N(t) = N^{1/2} \left(\frac{\tilde{c}_N(t)}{N} - \tilde{\xi}(t) \right)$$

converges weakly to a bivariate diffusion. Proceeding further, evaluate the characteristic function of the limiting noise,

$$E[\exp(i\theta_1 y_1(t) + i\theta_2 y_2(t))] = \phi(\theta_1, \theta_2, t); \quad (A-12)$$

The latter satisfies the following partial differential equation, from (2.7,B):

$$\begin{aligned}
\Phi' = & [-\theta_1 \lambda_{10} + (-\theta_1 + \theta_2) \lambda_{12}] \Phi_{\theta_1} - \theta_2 \lambda_{20} \Phi_{\theta_2} \\
& - \frac{1}{2} [\theta_1^2 \lambda_{01} + \theta_1^2 \lambda_{10} \xi_1(t) + (-\theta_1 + \theta_2)^2 \lambda_{12} \xi_1(t) + \theta_2^2 \lambda_{20} \xi_2(t)] \Phi
\end{aligned}
\tag{A-13}$$

This is recognizable as describing a bivariate Ornstein-Uhlenbeck process. Now one may differentiate (A-13) at $\theta_1 = \theta_2 = 0$, utilizing the fact that

$$V_i(t) = E[y_i^2(t)] = -\Phi_{\theta_i \theta_i}(0, 0, t), \quad i = 1, 2,$$

and

$$V_{12}(t) = E[y_1(t) y_2(t)] = -\Phi_{\theta_1 \theta_2}(0, 0, t)$$

to derive differential equations for the elements of the variance-covariance matrix of y . Not surprisingly, these differential equations agree with (23) (with λ_{ij} constant, and, in this case, $\lambda_{21} = 0$), i.e., $V_1(t) = \sigma_{11}(t)$, $V_{12}(t) = \sigma_{12}(t)$, $V_2 = \sigma_{22}(t)$. Thus if the same initial conditions are adopted for ξ, ζ and for y , and \tilde{x} the deterministic and noise components of the approximations of this paper and of B are seen to be identical. The validation can be extended with no difficulty in principle to the general situation.

2. Relation to Infinite-Server and Linear Markov Population Models

The class of compartment models described by the transition scheme (1) are similar to the classical Poisson-arrival infinite-server models of queueing theory and identical to certain Markov population models of Bartlett (1949), and Kingman (1969). We now briefly explore the connection, illustrating by the single-compartment and two-compartment examples. In particular, it will be shown that, even in the time-dependent parameter case, limiting Gaussian marginal distributions occur for compartment contents, in agreement with the development of the paper.

Single-Compartment Model. Let $C_1(t,u)$ denote the number of arrivals to the system that have occurred in the time interval (u,t) , $0 \leq u \leq t$, and that are present in compartment 1 at time t . Let $I_1(t,u)$ be the probability that an arrival occurring at time u is present in compartment 1 at time t . Consider the generating function of $C_1(t,u)$,

$$E[z_1^{C_1(t,u)}] = g(z_1, t; u) \quad (A-14)$$

Now by the assumption of Poissonian arrivals, see (1), we may write

$$g(z_1, t; u) = [z_1 I_1(t, u) N \lambda_{01}(u) du + 1 - I_1(t, u) N \lambda_{01}(u) du] g(z_1, t; u + du), \quad (A-15)$$

where the bracketted expression is the generating function of the contribution to $C_1(t, u)$ from arrivals during $(u, u + du)$; independence permits the multiplication. Subtract $g(z_1, t; u + du)$ from both sides, divide by du , and let $du \rightarrow 0$. The differential equation

$$-\frac{dg(z_1, t; u)}{du} = g(z_1, t; u) [(z_1 - 1) I_1(t, u) N \lambda_{01}(u) du] \quad (A-16)$$

results, the solution of which is, when u is set equal to zero,

$$g(z_1, t; 0) = E[z_1^{C_1(t)} | C_1(0) = 0] = \exp[(z_1 - 1) N \int_0^t I_1(t, u) \lambda_{01}(u) du], \quad (A-17)$$

the generating function of the Poisson distribution with mean $N \int_0^t I_1(t, u) \lambda_{01}(u) du$. It now follows immediately from the central limit theorem that the distribution of

$$x_1(t) = \frac{C_1(t) - N \int_0^t I_1(t, u) \lambda_{01}(u) du}{\sqrt{N}} \quad (A-18)$$

converges weakly to the normal law with mean zero and variance $\int_0^t I_1(t, u) \lambda_{01}(u) du$. Furthermore, it is easy to verify that

$$\text{Var}[y(t)] = \int_0^t I_1(t, u) \lambda_{01} du = \frac{\lambda_{01}}{\lambda_{10}} (1 - e^{-\lambda_{10} t})$$

when λ_{01} and λ_{10} are both constants, and $y(t)$ is the solution of the stochastic differential equation (A-7) which results from following the approximation program that is the topic of the paper; we start with $C_1(0) = 0$. Verification in the time-dependent parameter case is also possible, but is more difficult. Thus the correct marginal distribution is verifiable directly for this, the simplest, case. With more labor the joint distribution at different times may be found using an extension of the technique.

Two-Compartment Model. Consider the generating function of $(C_1(t,u), C_2(t,u))$, denoted by $g(z_1, z_2, t; u)$. Then an argument analogous to that joint presented shows that g satisfies a differential equation, which when solved yields

$$g(z_1, z_2, t; 0) = \exp \left\{ N \left\{ (z_1 - 1) \int_0^t I_1(t, u) \lambda_{01}(u) du + (z_2 - 1) \int_0^t I_2(t, u) \lambda_{01}(u) du \right\} \right\} \quad (A-19)$$

implying that the contents of the two compartments are Poisson distributed at time t , and that $C_1(t)$ and $C_2(t)$ are independent. The same central limit theorem argument now shows joint normality as $N \rightarrow \infty$ in agreement with the claims of the body of the paper. Calculations omitted here show that the diffusion approximation produces first and second moments that agree with those of the Poisson input model above. In particular, the correlation term, σ_{12} , of (20) equals zero, signifying the independence that we noted in (A-19).

Distribution List

| | Copies |
|---|--------|
| Defense Documentation Center Cameron Station Alexandria, Virginia 22314 | 2 |
| Dean of Research Code 023 Naval Postgraduate School Monterey, California 93940 | 1 |
| Library (Code 0212) Naval Postgraduate School Monterey, California 93940 | 2 |
| Library (Code 55) Naval Postgraduate School Monterey, California 93940 | 2 |
| D. R. Cox Department of Mathematics Imperial College London, SW7 England | 1 |
| D. R. McNeil Department of Statistics Princeton University Princeton, New Jersey 08540 | 1 |
| J. A. Hooke E. Wolman Bell Telephone Labs Holmdel, New Jersey 07733 | 1 1 |
| B. J. McDonald Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 | 1 |
| L. Kleinrock R. Muntz Computer Science University of California Los Angeles, California 90024 | 1 1 |
| J. M. Wozencraft Electrical Engineering M.I.T. Boston, Massachusetts 02139 | 1 |

| | Copies |
|---|--------|
| D. L. Iglehart Department of Operations Research Stanford University Stanford, California 94305 | 1 |
| J. Riordan Department of Mathematics Rockefeller University New York, New York 10021 | 1 |
| G. Fishman University of North Carolina Chapel Hill, North Carolina 27514 | 1 |
| P. T. Holmes Department of Mathematics Clemson University Clemson, South Carolina 29631 | 1 |
| R. Elashoff Biomathematics University of California Los Angeles, California 90024 | 1 |
| Gene Fisher Management Science Department RAND Corporation 1700 Main Street Santa Monica, California 90401 | 1 |
| R. M. Stark Statistics and Computer Science University of Delaware Newark, Delaware 19711 | 1 |
| M. Mazumdar Robert Hooke Mathematics Department Westinghouse Research Labs Churchill Boro Pittsburgh, Pennsylvania 15235 | 1 1 |
| J. R. Capra Center for Naval Analysis 1401 Wilson Boulevard Arlington, Virginia 22209 | 1 |
| Roy Welsch Sloan School M.I.T. Cambridge, Massachusetts 02139 | 1 |

| | Copies |
|---|--------|
| J. R. Blum | 1 |
| B. R. Agins | 1 |
| Mathematical Science National Science Foundation 18th and "G", N.W. Washington, D.C. | |
| Dr. A. J. Goldman | 1 |
| Chief, Operations Research 205.02 Admin., A428 National Bureau of Standards Washington, D.C. 20234 | 1 |
| Dr. Leon F. McGinnis | 1 |
| School of Ind. and Systems Eng. Georgia Institute of Technology Atlanta, Georgia 30332 | |
| Dr. H. Higgins | 1 |
| 53 Bonn 1, Postfach 589 Nassestrasse 2 West Germany | |
| Prof. Saul I. Gass | 1 |
| College of Bus. and Management University of Maryland College Park, Maryland 20742 | |
| Dr. V. Srinivasan | 1 |
| Graduate School of Business Stanford University Stanford, California 94305 | |
| Code 303 Sorenson | 1 |
| Navy Per. Res. and Dev. Ctr. 271 Catalina Blvd. San Diego, California 92152 | |
| Dr. Paul Schweitzer | 1 |
| Thomas J. Watson Research Ctr. Post Office Box 218 Yorktown Heights New York 10598 | |
| Dr. Linus Schrage | 1 |
| University of Chicago Graduate School of Business 5836 Greenwood Avenue Chicago, Illinois 60637 | |
| For Math. Studies in E Ctr. and Man. Sci. Northwestern University Evanston, Illinois 60201 | 1 |

| | Copies |
|---|--------|
| Sci. Res. Ctr. Man. Faculty of Com. and Bus. Admin. University of British Columbia Vancouver British Columbia V6T 1W5 Canada | 1 |
| Operations Research Attn: T. Varley, Code 434 Office of Naval Research Arlington, Virginia 22217 | 1 |
| John Lehoczky Statistics Department Carnegie-Mellon University Pittsburgh, Pennsylvania 15213 | 1 |
| D. P. Gaver Code 55Gv Naval Postgraduate School Monterey, California 93940 | 25 |
| H. Kobayashi M. Reiser IBM Thomas J. Watson Research Center Yorktown Heights, New York 10598 | 1 1 |
| R. Van Slyke Network Analysis Corporation Beechwood, Old Tappan Road Glen Cove, New York 11542 | 1 |